

SPATIAL DIFFUSION OF X-RAY EMISSION LINES IN THE M87 COOLING FLOW; EVIDENCE FOR ABSORPTION

WILLIAM G. MATHEWS¹, DAVID A. BUOTE^{1,2,4}, AND FABRIZIO BRIGHENTI^{1,3}

¹University of California Observatories/Lick Observatory, Board of Studies in Astronomy and Astrophysics,
 University of California, Santa Cruz, CA 95064
 mathews@ucolick.org

²Department of Physics and Astronomy, University of California, Irvine, CA 92697
 buote@uci.edu

³Dipartimento di Astronomia, Università di Bologna, via Ranzani 1, Bologna 40127, Italy
 brighenti@bo.astro.it

Draft version February 1, 2008

ABSTRACT

Recent XMM-Newton observations of the cooling flow in M87 indicate sharply decreasing oxygen, iron and silicon abundances within ~ 5 kpc of the galactic center. This result is unexpected since stellar mass loss and Type Ia supernovae are expected to produce pronounced central abundance maxima for all three elements. However, it has been suggested that many of the strong X-ray lines are optically thick and diffuse to larger radii in the cooling flow before escaping, falsifying the central abundances. We verify with radiation transfer calculations that this effect does indeed occur in the M87 cooling flow, but that it is insufficient to account for the M87 observations. We suggest that some source of continuous opacity is required to reduce the central X-ray line emission, perhaps by warm gas at $T \sim 10^5 - 10^6$ K. The radial surface brightness profiles of X-ray resonance lines are also sensitive to turbulence in cooling flows which reduces the line center optical depths considerably. Turbulence may provide sufficient energy to continuously heat the warm absorbing gas.

Subject headings: galaxies: elliptical and lenticular – galaxies: cooling flows – galaxies: interstellar medium – X-rays: galaxies – X-rays: galaxy clusters

1. DIFFUSION OF X-RAY LINE PHOTONS IN M87

This estimate of the diffusion of X-ray line photons in M87 has been motivated by the exceptionally low abundances of iron, silicon and oxygen within ~ 5 kpc of the center of M87 as recently observed by Böhringer et al (2000) with XMM-Newton. We explore the possibility that these abundance variations are artifacts of the outward diffusion of optically thick X-ray lines as suggested by Gil'franov et al. (1987), Tawara et al. (1997), Shigeyama (1998) and Böhringer et al. (2000). We find that the expected outward migration of line photons does occur, but is insufficient to account for the central photon depletion observed. Instead, an additional source of continuous absorption is indicated, as suggested by Buote (2000), and we speculate about its physical nature. Before discussing our results, we begin with a review of the properties of the M87 cooling flow and describe our line transfer calculation.

2. VARIATION OF GAS TEMPERATURE AND DENSITY

The electron density distribution in the M87 cooling flow is taken from Nulsen & Böhringer (1995), corrected to distance 17 Mpc ($1' = 4.945$ kpc):

$$n_{e,g} = n_0[1 + (r/r_o)^p]^{-1} \quad (1)$$

with $n_0 = 0.120 \text{ cm}^{-3}$, $p = 1.18$ and $r_o = 0.5r_e$ where $r_e = 8.625$ kpc is the effective radius. According to Böhringer et al. (2000) these observations underestimate the central density which should instead be represented by a β model fit:

$$n_{e,\beta} = n_1[1 + (r/r_\beta)^2]^{-3\beta/2} \quad (2)$$

with $n_1 = 0.35 \text{ cm}^{-3}$, $\beta = 0.47$ and $r_\beta = 1.97$ kpc. This inner density distribution is based on ROSAT HRI observations reported in Böhringer (1999). Since the central X-ray emission in M87 is known to be partly non-thermal (and time variable), this latter density determination may be uncertain. Nevertheless, in the following we assume that $n_e(r)$ is given by the maximum of $n_{e,g}(r)$ and $n_{e,\beta}(r)$ which intersect at 11.9 kpc.

The (single-phase) gas temperature distribution is found by combining the results of Böhringer et al. (2000) within 65 kpc with more distant temperatures measured by Nulsen & Böhringer (1995). These results can be fit with the fitting function:

$$T = T_o \left(\frac{r_m}{r_{ot}} \right) \left[\frac{r_m}{(r + r_{ot})} + \left(\frac{r}{r_m} \right)^q \right]^{-1} \quad (3)$$

with $T_o = 1.5 \times 10^7$ K, $r_m = 5.695$, $r_{ot} = 1.53$ and $q = 0.22$; all radii are in units of r_e . This temperature profile rises from the center toward a broad maximum, similar to thermal profiles in other well-observed bright elliptical galaxy cooling flows (Brighenti & Mathews 1997). The location of the temperature maximum in M87 is at an unusually large radius presumably due to the relatively high ambient temperature (2 - 3 keV) in the Virgo cluster. The central cooling of the gas can be understood as the mixing of hot inflowing gas from the Virgo cluster with gas lost from evolving stars in M87 which orbit with a lower virial temperature than that of the surrounding cluster (Brighenti & Mathews 1999a). The characteristic deep central temperature minimum in M87 indicates that most of the gas in M87 has not been substantially heated by the jet and

⁴ Chandra Fellow.

radio source.

3. ABUNDANCES, LINE EMISSION AND ABSORPTION

We assume meteoritic solar abundances with $\log(a_E) \equiv \log(n_E/n_H) = -3.07$ and -4.45 for O and Si respectively. The radial abundance variation of each element in the hot gas relative to solar meteoritic is represented with $A_E(r)$. The line emissivity is assumed to vary with local temperature according to the emission coefficients Λ_ℓ determined by Landini & Monsignori Fossi (L-MF; 1990):

$$\varepsilon_\ell = A_E(r)n_en_H\alpha\Lambda_\ell(T) \quad (\text{erg cm}^{-3} \text{ s}^{-1})$$

where α is a factor of order unity that corrects L-MF abundances to solar meteoritic and $n_H = n_e(4 - 3\mu)/(2 + \mu)$ is the local proton density in a plasma with molecular weight $\mu = 0.61$.

The emission lines that we consider arise from the ground state and are Doppler broadened by thermal and turbulent velocities. The absorption coefficient is

$$\kappa(x) = n_H x_i a_E A_E \left(\frac{\pi e^2}{m_e c} f_{gu} \frac{1}{\pi^{1/2}} \frac{1}{\Delta\nu_D} \right) e^{-x^2} \quad (\text{cm}^{-1}).$$

Here $a_E A_E(r)$ is the local abundance of the line-emitting element and $x_i(r)$ is the fractional abundance of the line-emitting ion interpolated from the results of Sutherland & Dopita (1993) for collisional ionization equilibrium. Photons are assumed to be emitted in a Gaussian profile centered on the line energy $E_o = h\nu_o$ and

$$x = \frac{\nu - \nu_o}{\Delta\nu_D}$$

is the dimensionless line frequency with

$$\Delta\nu_D = \frac{\nu_o}{c} \left(\frac{2kT}{A_{Ewt}m_p} + v_{turb}^2 \right)^{1/2} \quad (4)$$

where A_{Ewt} is the atomic weight of element E. The line center optical depth in the radial direction is found by integrating $d\tau_o = \kappa(0)dr$.

We are taking $x_i(T)$ and $\Lambda_\ell(T)$ from two different sources. However, the error from this possible inconsistency is unlikely to be large since collisional ionization fractions $x_i(T)$ found by most authors, including L-MF, are in good agreement with those of Arnaud & Rothenflug (1985).

4. MONTE CARLO PROCEDURE

To estimate the diffusion of X-ray line photons, we represent the inner cooling flow of M87 with $N = 50$ radial zones, each having different but individually uniform temperature and density. The zone spacing r_n is chosen so that the line luminosity is the same from each zone. The Monte Carlo process begins by choosing the zone of origin of each photon based on equal weighting among the zones. The exact birth radius r_b of each photon within the chosen zone is then found from

$$r_b = [r_n^3 + \mathcal{R}(r_{n+1}^3 - r_n^3)]^{1/3}$$

where $\mathcal{R}$ is a random number between 0 and 1. The initial direction of the photon is $w = \cos\theta = 2\mathcal{R} - 1$ where $\mathcal{R}$ is another random number and θ is the angle between the photon velocity and the direction to the center of the cooling flow. The initial frequency x is chosen by random

selection from a Gaussian probability density. The distance that the photon moves from its origin r_o to the local zone boundary r_z is

$$\ell = wr_o + \sigma[r_z^2 - r_o^2(1 - w^2)]^{1/2}. \quad (5)$$

Here $\sigma = 1$ and $r_z = r_{n+1}$ for outward moving photons ($w < w_{cr}$) and $\sigma = -1$ and $r_z = r_n$ for inward moving photons ($w > w_{cr}$) where $w_{cr} = [1 - (r_n/r_o)^2]^{1/2}$. In the zone of photon birth $r_o = r_b$. If the photon continues to move through several zones without scattering, w and r_o remain constant and Equation (5) can be used to find the total path length moved L . The distance ℓ traversed by the photon across each zone must be compared to the free path in the zone:

$$\ell_s = -\ln(\mathcal{R})/\kappa_n(x_n)$$

where $\kappa_n(x_n)$ is the absorption coefficient of the photon in zone n . Since $\Delta\nu_D$ varies from zone to zone it is necessary to rescale the frequency to the local zone n ($r_n < r < r_{n+1}$): $x_n(r) = x_n(r_b)[\Delta\nu_D(r_b)/\Delta\nu_D(r)]^{1/2}$. If $\ell_s > \ell$, the photon moves into the next zone, but if $\ell_s < \ell$ the photon is scattered at radius

$$r_s = (r_o^2 + L_s^2 - 2r_o L_s w)^{1/2}$$

where $L_s = \sum \ell_n + \ell_s$. The photon is then scattered into a new random direction beginning a new trajectory with $r_o = r_s$, assuming complete redistribution in the Gaussian line profile. The scattering is conservative so all photons must ultimately flow out beyond the outermost zone. When the photon reaches the outer boundary of the gas $r_t = r_N$, where the optical depths are very small, it is assumed to escape freely.

5. RESULTS AND DISCUSSION

We describe transfer calculations for two lines: SiXIV 2.003 keV ($L\alpha$) and OVIII 0.652 keV ($L\alpha$), both having $f_{gu} = 0.420$. Since these resonance lines are among the most optically thick lines of Si and O at M87 gas temperatures, the apparent abundance distribution inferred from each of these lines – subsequent to spatial diffusion – must correspond to the largest possible central abundance depletion. For each line-emitting element the apparent radial abundance profiles observed with XMM-Newton can be fit with

$$A_E^{(ob)}(r) = A_{ob} \left(\frac{r_{mob}}{r_{otob}} \right) \left[\frac{r_{mob}}{(r + r_{otob})} + \left(\frac{r}{r_{mob}} \right)^{qob} \right]^{-1}. \quad (6)$$

The apparent (meteoritic) abundance dependence of Si and O are approximated with: $(A_{ob}, r_{mob}, r_{otob}, qob) = (0.1, 7.4, 0.34, 0.7)$ and $(0.05, 8.0, 0.5, 0.94)$ respectively; r_{mob} and r_{otob} are in kpc. The line center optical depths in the three lines evaluated with $A_E^{(ob)}(r)$ (Fig. 1) are smaller than those suggested by Böhringer et al (2000) in part because we use the higher temperatures observed instead of their isothermal approximation. If the observed abundance maxima are due to outward photon diffusion by conservative scattering before escape, the total X-ray line luminosity within some larger radius (where $\tau \ll 1$) is unaffected by the radiative transfer.

According to this line diffusion hypothesis, the true radial abundance profiles are expected to decrease monotonically as we represent with

$$A_E(r) = \frac{A_o}{1 + (r/r_E)^c} + A_{cf}. \quad (7)$$

The first term accounts for the local enrichment by stellar mass loss and SNIa explosions, and the second term represents the abundance of cluster cooling flow gas flowing into M87, but the parameterized fit is not intended to exactly quantify these two enrichment sources. The coefficients in Equation (7) are chosen so that the total line luminosities within 50 kpc are very close to luminosities based on $A_E^{(ob)}$ (see below): $(A_o, r_E(\text{kpc}), c, A_{cf}) = (1.0, 35, 1.4, 0.15)$ and $(0.4, 12, 2, 0.14)$ for Si and O respectively. The A_{cf} are chosen for agreement with the XMM-Newton abundances at $r \gtrsim 30$ kpc; r_E is assumed to be similar to $r_e = 8.625$ kpc appropriate for a stellar contribution; c is chosen for a reasonable fit to the XMM-Newton data in $10 \lesssim r \lesssim 30$ kpc; A_o is adjusted until the total luminosity within $r = 50$ kpc (where all lines are optically thin) matches that of each line computed with $A_E^{(ob)}(r)$. With these assumed intrinsic abundance profiles $A_E(r)$, the total line luminosities within 50 kpc exceed the approximate observed luminosities of the Si and O lines by about 2 and 13 percent respectively. We exceed the oxygen line luminosity based on $A_O^{(ob)}(r)$ since this abundance is less accurately determined by the XMM-Newton data. The cumulative line center optical depths are somewhat greater when evaluated with $A_E(r)$ than with $A_E^{(ob)}(r)$ as shown in Figure 1.

Monte Carlo radiative transfer for each line has been calculated with 10^4 photons, 50 spatial zones and abundances $A_E(r)$. Results are illustrated in Figure 2. The light dotted histograms show the abundance of each element that would be inferred by observing each line in the optically thin limit, $A_E(\text{birth}) = A_E(r)N_{ph,b}/\langle N \rangle$, where $N_{ph,b}$ is the number of photons created in each zone and $\langle N \rangle = 200$ is the mean number of photons per zone. The adjacent light-line histograms show the variation of the apparent abundance indicated by each line after radiative diffusion, $A_E(\text{escape}) = A_E(r)N_{ph,e}/\langle N \rangle$, where $N_{ph,e}$ is the number of photons that escape from each zone with no further scattering. $A_E(\text{escape})$ can be compared directly with the observed abundances.

The light solid-line histograms clearly indicate central photon depletions or flattening within ~ 5 kpc. The apparent Si abundance based on the 2.003 keV line is clearly self-absorbed and matches the observations quite well at $r \gtrsim 3$ kpc. But the amount of photon depletion with conservative scattering is clearly insufficient to explain the observed O abundance. It is not possible to deepen the central minima by increasing the O abundance $A_O(r)$ (and line opacity) in $r \lesssim 5$ kpc since the global line luminosities would exceed those observed. The OVIII line luminosity may already be too high by about 13 percent. The central oxygen line luminosity (abundance) only levels off and does not decrease toward the origin as observed. Evidently line photons are being lost by absorption. The central abundances of Si and O – $A_E(0) = 1.15$ and 0.54 respectively – cannot be trusted if the line scattering is non-conservative. Since the continuous opacity in a warm plasma with $T_w \lesssim 10^6$ K decreases rapidly with photon energy, $\sigma \propto E^{-3}$, the OVIII line should suffer ~ 30 times more absorption than the SiXIV line, in qualitative agreement with the profiles in Figure 2.

Shigeyama (1998) noted that if the outward diffusion of

strong X-ray lines is not considered in the data reduction, the central gas density can be underestimated by $\sim 20 - 50$ percent. To explore this, we repeated the calculations with $n_{e,\beta}$ increased by 1.5. Our conclusions are strengthened by this adjustment in the gas density. With higher $n_{e,\beta}$ the luminosity of the lines within 50 kpc can be matched with a slightly lower central abundance, resulting in central minima in $A_E(\text{escape})(r)$ even less pronounced than those in Figure 2.

Studies of optical line emission in the central regions of galactic and galaxy group cooling flows reveal widespread random motions typically about 0.2 – 0.4 of the sound speed in the hot gas (Heckman et al. 1989; Caon et al. 2000). Since small, optically visible line-emitting regions at $T \sim 10^4$ K are likely to be strongly coupled to the ambient hot gas, we infer that the hot gas must share the same turbulent velocities. Turbulence is expected to strongly influence the spatial distribution of X-ray resonance lines, particularly for more massive line-emitting ions (Equation 4). To demonstrate this, we repeated the transfer calculations for both lines with $v_{turb}^2 = \mathcal{M}_{turb}^2(5A_E/6\mu)$ assuming a turbulent Mach number $\mathcal{M}_{turb} = 0.3$. In the presence of turbulence, which we regard as more realistic, the line center optical depths are greatly reduced (Fig. 1), there is less outward diffusion (Fig. 2), and the apparent central depletion of silicon photons is significantly reduced.

We have also computed line profiles for the FeXXV 6.69 keV (He 4) resonance line discussed by Böhringer et al. but have not discussed its profile here because emission in this line is unlikely to be representative of the intrinsic iron abundance. The FeXXV 6.69 keV line is very sensitive to the (low) cooling flow temperatures $T(r)$ in $r \lesssim 10$ kpc where the line transfer is most relevant (e.g. Fig. 2 of Buote, Canizares & Fabian 1999) and relatively few FeXXV line photons are created in this low temperature region. Because of this, the χ^2 iron abundance determination in M87 by Böhringer et al. is dominated by the multitude of strong Fe L lines near ~ 1 keV. The many lines that constitute the Fe L feature have different atomic parameters, so no single L line can faithfully represent the iron abundance, as we have assumed for the O and Si Lyman- α lines. Since the continuous plasma opacity varies as $\sigma \propto E^{-3}$, the opacity at the FeXXV 6.69 keV is ~ 1000 times less than the opacity at the OVIII 0.652 keV line. An observation of the equivalent width of the essentially unabsorbed FeXXV 6.69 keV line with projected radius in M87 would allow a calculation of the intrinsic iron abundance distribution, provided $T(r)$ is sufficiently well determined.

We conclude that the central minima in apparent abundances and line luminosities in M87 cannot be understood solely in terms of conservative photon scattering. An additional source of continuous opacity seems to be required. We speculate that the source of this opacity is the warm absorbing gas at $T \sim 10^5 - 10^6$ K as suggested for M87 by Buote (2000). Buote's conclusion is derived from modeling the ROSAT PSPC data of M87 with coronal plasma emission modified by a single absorption edge and Galactic absorption. The rest energy of the edge is determined to lie at ~ 0.6 keV, near the position of the OVIII L α line. The absorption derived from the ROSAT data ($\tau \sim 0.5 - 1$) is spatially more extended than that required to account for

central minima in the X-ray resonance lines. Because of the limited energy resolution of ROSAT, the effect of using a simple absorption edge cannot be distinguished from a model in which the O abundance is allowed to take low values as done by Böhringer et al (2000).

The central declines in Fe and Si abundances in the Böhringer et al. data are difficult to understand since we expect hot gas closer to the galactic core to be more enriched by centrally peaked stellar mass loss and Type Ia supernovae (Brighenti & Mathews 1999b). Since the oxygen abundance found by Böhringer et al. has a generally negative gradient beyond $\sim r_e$, it is apparent that the stellar and Type Ia ejecta are more oxygen-rich than cooling flow gas entering M87 from larger radii, so a central peak in the O abundance would also be expected. The need for an additional source of continuous opacity in our line transfer calculations lends new support to the absorption interpretation. If this absorption arises from warm gas, then its mass is similar to that of the hot gas (Buote 2000).

The energy source that heats the warm gas is of great interest. While some of the cooling flow gas may be heated by the M87 jet, this region is avoided in the Böhringer et al. observations; the deep central minimum in their $T(r)$ shows no evidence of AGN heating. The most likely energy source for heating the warm gas is the turbulent motion of the hot gas as indicated by the dispersion velocity of

optical emission lines $v_t \sim 200 \text{ km s}^{-1}$ in M87 (Heckman et al. 1989). These velocities are subsonic in the hot gas, but would be supersonic in the warm, X-ray absorbing gas. Colliding regions of warm gas would be shock-heated at a rate $H = f n_c m_p v_t^3 / r_c \text{ ergs cm}^{-3} \text{ s}^{-1}$ where f is the filling factor of warm gas clouds of mean radius r_c and electron density n_c . Although we expect that some net cooling occurs, if shock heating balances radiative cooling $C = n_c^2 \Lambda \text{ ergs cm}^{-3} \text{ s}^{-1}$ then the cloud size can be estimated: $r_c \sim f m_p v_t^3 / n_c \Lambda \sim 400 f \text{ pc}$ for $v_t = 200 \text{ km s}^{-1}$, $n_c = 0.1 \text{ cm}^{-3}$ and cooling coefficient $\Lambda = 10^{-22} \text{ erg cm}^3 \text{ s}^{-1}$. With $f \sim 0.5$, this cloud size is consistent with the scale of irregularities observed in the optical line emission and the relative uniformity of the warm plasma absorption.

Finally, we speculate further that such extended absorbing regions are generally present in cooling flows and account for the apparent deficiency of emission lines characteristic of cooling ions in the centers of these flows (e.g. Peterson et al. 2000; McNamara et al. 2000).


We thank referee Paul Nulsen for his helpful suggestions. Studies of the evolution of hot gas in elliptical galaxies at UC Santa Cruz are supported by NASA grant NAG 5-3060 and NSF grant AST-9802994 for which we are very grateful. DAB was supported by NASA Chandra Fellowship grant PF8-10001 awarded by the Smithsonian Astrophysical Observatory under contract NAS8-39073.

REFERENCES

- Anders, E. & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
 Arnaud, M. & Rothenflug, R. 1985, *A&AS*, 60, 425
 Böhringer, H. 1999, in *Diffuse Thermal and Relativistic Plasma in Galaxy Clusters*, H. Böhringer, L. Feretti, P. Schuecker (eds.) MPE Report, p. 115
 Böhringer, H. et al. 2000 *A&A* (submitted) (astro-ph/0011459)
 Brighenti, F. & Mathews, W. G. 1997, *ApJ*, 486, L83
 Brighenti, F. & Mathews, W. G. 1999a, *ApJ*, 512, 65
 Brighenti, F. & Mathews, W. G. 1999b, *ApJ*, 515, 542
 Buote, D. A. 2000, *ApJ* (in press) (astro-ph/0008408)
 Buote, D. A., Canizares, C. R. & Fabian, A. C. 1999, *MNRAS* 310, 483
 Caon, N., Macchetto, D. & Pastoriza, M. 2000, *ApJS* 127, 39
 Gil'fanov, M. R., Sunyaev, R. A. & Churazov, E. M. 1987, *Sov. Astron. Lett.*, 13, 3 (*Pis'ma Astron. Zh.* 13, 7, 1987)
 Heckman, T., Baum, S. A., van Breugel, W. J. M. & McCarthy, P. 1989, *ApJ*, 338, 48
 Landini, M. & Monsignori Fossi, B. C. 1990, *A&AS* 82, 229
 McNamara, B. R. et al. (2000) in "Constructing the Universe with Clusters of Galaxies," IAP, Paris (in press) (astro-ph/0012331)
 Nulsen, P. E. J. & Böhringer, H. 1995, *MNRAS* 274, 1093
 Peterson, J. R. et al. (2000) *A&A* (in press) (astro-ph/0010658)
 Shigeyama, T. 1998, *ApJ*, 497, 587
 Sutherland, R. S. & Dopita, M. A. 1993, *ApJS*, 88, 253
 Tawara, Y. et al. 1997, in *X-ray Imaging and Spectroscopy of Cosmic Hot Plasmas*, F. Makino & Mitsuda (eds.), Universal Academy Press, Tokyo, p. 87

FIG. 1.— Variation of line center optical depths in the central cooling flow of M87 in two lines: SiXIV 2.003 keV and OVIII 0.652 keV. The solid and dashed lines show optical depths resulting from the model $A_E(r)$ and directly from the observations $A_E^{(ob)}(r)$ respectively. The heavy dashed lines are line center optical depths based on $A_E(r)$ but in the presence of turbulence with $M_{turb} = 0.3$.

FIG. 2.— Radial variation of silicon and oxygen abundances in the M87 cooling flow. The observed points including symbol type have been taken directly from Böhringer et al. (2000). The smooth lines passing through the observations are $A_E^{(ob)}(r)$ found from Equation (6). The dotted histograms show the variation of the elemental abundances $A_E(birth)$ based on each line using the Monte Carlo program in the optically thin limit; apart from statistical fluctuations this histogram is identical to $A_E(r)$. The light solid histograms are the abundance profiles $A_E(escape)$ that would be inferred from each resonance line after Monte Carlo spatial diffusion; the line center optical depths for these calculations are the light solid lines from Figure 1. The heavy histograms show the effect of turbulence with $M_{turb} = 0.3$.



$A_E(\text{birth})$ and $A_E(\text{escape})$

